

A projection operator technique for solution of relativistic wave equation on non-compact group: the case of a charged vector-boson

Halina V. Grushevskaya¹ and Leonid I. Gurskii²

¹ *Physics Department, Belarusian State University,
4 Fr. Skorina Av., 220050 Minsk, Belarus*

² *Belarusian State University of Informatics and Radio electronics,
6 P. Brovka Str, 220027 Minsk, Belarus*

E-mail: grushevskaja@bsu.by

Abstract

A projection operator technique for solution of relativistic wave equation on non-compact group has been proposed. This technique was applied to the construction of wave equations for charged vector boson in a potential field. The equations were shown to approximately describe a hydrogen-like atom and allow estimating of relativistic corrections such as a fine structure of hydrogen atom lines with high accuracy.

PACS numbers: 02.30.Tb; 03.65.Pm; 02.40.k

Keywords: projection operator, charged vector-boson, non-compact group, composite quantum system

1 Introduction

Geometrical approach plays an important role in the study of quantum models and relativistic processes [1, 2, 3]. This paper discusses the projection operator technique to examine motion of a relativistic vector boson with a non-compact symmetry group $SO(4,2)$. The constructed equations of motion can also be used for the description of a hydrogen-like atom, since being a composite quantum system, it has an integer-valued spin. In the paper, these equations are shown to estimate relativistic corrections such as a fine structure of hydrogen atom lines with high accuracy.

The goal of the paper is to develop a geometrical method of the projection operators and on this base to examine motion of charged vector boson in a potential field.

2 Projection operator technique

Let us consider an equation for a relativistic quantum particle in two-dimensional flat space with two complex coordinates

$$|\xi_s\rangle, |\xi_{\dot{s}}\rangle, \quad s(\dot{s}) = 1, 2; \quad (1)$$

being components of ket bispinor.

Consider motion of a free particle, then the Hamiltonian $\mathcal{H}$ includes only relativistic kinetic energy term T . For comparison, we shall remind a classic analog $T^2 = Z^2 + \vec{k}^2$. Here $\vec{k}$ is a momentum, Z is a "mass", c is the velocity of light: $c = 1$. Then, a relativistic wave equation reads

$$T|\psi\rangle = i\frac{\partial|\psi\rangle}{\partial t} \quad (2)$$

where $|\psi\rangle$ is a wave function, t is time, i is imaginary unit, $\hbar = 1$. Eq. (2) can be rewritten as

$$\left(\mathcal{H}^\dagger \mathcal{H} + \frac{\partial^2}{\partial t^2}\right)|\psi\rangle = \left(-\frac{\partial^2}{\langle\partial\xi_s|\partial\xi_s\rangle} - Z^2 + \frac{\partial^2}{\partial t^2}|\xi_s\rangle\langle\xi_s|\right)|\psi\rangle \equiv (H - Z^2)|\psi\rangle = 0. \quad (3)$$

Here the summation is assumed over repeated indexes, and it has been taken into account that the following expression holds: $|\xi_s\rangle\langle\xi_s| = \hat{I}$, where $\hat{I}$ is the operator unity; the wave function $\langle\xi_s|\psi\rangle$ is considered in the representation $|\xi_s\rangle$, $s = 1, 2$. It follows from Eq. (3) that it is possible to consider the square of Z as eigenvalues of the operator H , which is equal to $i\frac{\partial}{\partial t_1}$, where t_1 is a time coordinate which does not depend on t . Hence, the equation (3) is determined on the manifold with four space-like and two time-like coordinates.

Let us pass into the representation $\{|\phi_i\rangle\}$ where the operator representation $\hat{\xi} \equiv T_\xi$ of a translation group $\{\hat{\xi}\}$ has the following form: $\hat{\xi} \rightarrow i\frac{\partial}{\partial k_i}$, k_i is a momentum. Then, using the properties of the projection operators [4] Eq. (3) can be rewritten as

$$\left(-\frac{\partial^2}{\langle\partial\xi_s|\partial\xi_s\rangle} - Z^2 + \frac{\partial^2}{\partial t^2}\delta_{ki}(|\phi\rangle\langle\phi_k|\xi_s\rangle\langle\xi_s|\phi_i\rangle\langle\phi_r|)\right)|\psi_r\rangle = 0. \quad (4)$$

The components of the wave function for the free particle in this representation are

$$\langle \phi_i | \psi \rangle = \exp \left\{ i \left(k_s \xi_{si} + \epsilon_{srq} k_s \frac{1}{|\xi|^2} [\xi_{ri}, \xi_{qi}] \right) \right\} \exp(i\omega t), \quad (5)$$

$$\xi_{si} \equiv \langle \phi_i | \xi_s \rangle. \quad (6)$$

Here the first term of the expression (6) describes the translation to the group unity; the commutator appeared in the second summand is an element of the rotation group; ω is a frequency, $|\xi|^2 = \sum_{\nu=1}^4 \xi_\nu^2$, ξ_{si} are components of bispinor in representation $\{\phi_i\}_{i=1}^N$, $N = 1, 2, \dots$, being c -numbers; ϵ_{ijk} is the Levi-Civita tensor.

Using the definition of a projector, after differentiation with respect to $t, \vec{k}$, we transform the equation (4) to the following form

$$\left(-\frac{\partial^2}{\partial \xi_s \partial \xi_s} - Z^2 + \omega^2 (\xi_{si} + \epsilon_{srq} [\xi_{ri}, \xi_{qi}]) (\xi_{si} + \epsilon_{skq} [\xi_{ki}, \xi_{qi}]) |\phi_i\rangle \langle \phi_r| \right) \psi_r = 0 \quad (7)$$

Assuming that the basis $\{\xi_{si}\}$ obeys a condition of orthogonality, finally Eq. (7) is written in the reduced form

$$\left(-\frac{\partial^2}{\partial \vec{\xi}_s \partial \vec{\xi}_s} - Z^2 + \omega^2 \left(\vec{\xi}_s \vec{\xi}_s + \epsilon_{srq} \frac{1}{|\xi|^4} [\vec{\xi}_r, \vec{\xi}_q] \epsilon_{srq} [\vec{\xi}_r, \vec{\xi}_q] \right) |\Phi\rangle \langle \Phi| \right) \psi = 0 \quad (8)$$

where $\vec{\xi}_s \stackrel{def}{=} \{\xi_{si}\}_i$ are the components of bispinor in the basic set $\Phi \stackrel{def}{=} \{|\phi_i\rangle\}$.

To find a real equation of motion, let us consider particle motion in three-dimensional space. It is known [5] that the mapping of three-dimensional sphere $S^3 \in \mathbb{R}^4$ into three-dimensional space $\mathbb{R}^3$ is two-valued. Therefore, the set of four numbers describing a position of a particle in four-dimensional space, is replaced by a set of four numbers $\{x_\lambda, S\}$, $\lambda = 1, 2, 3$ also; but one of them (S) is a discrete quantity. S takes two values $S = \pm 1$ depending on either we exclude north or south pole when projecting. The physical interpretation of S is the helicity of a particle as a result of the projection P on three-dimensional subspace. For the ket vector $|x_\lambda^{R(L)}\rangle$ the symbols R, L are attributed to right- and left-helical particles, respectively. Then, we can write the following identity

$$\begin{aligned} & \sum_\mu |\xi_\mu\rangle \langle \xi_\mu | \psi \rangle + \frac{1}{2} \sum_\lambda [|\xi_\kappa\rangle (\sigma_\lambda^+)_{\kappa\nu} \langle \xi_\nu | \psi \rangle + |\xi_\kappa\rangle (\sigma_\lambda^-)_{\kappa\nu} \langle \xi_\nu | \psi \rangle] \\ & \equiv \sum_\lambda \left[\frac{1}{2} (|x_\lambda^L\rangle \langle x_\lambda^L| + |x_\lambda^R\rangle \langle x_\lambda^R|) \psi + \frac{1}{2} (|x_\lambda^L\rangle \langle x_\lambda^L| + |x_\lambda^R\rangle \langle x_\lambda^R|) \psi \right], \end{aligned} \quad (9)$$

where matrices $\sigma_\lambda^\pm$ are determined by

$$\sigma_\lambda^+ = \epsilon_{\lambda ij} \sigma_{ij}; \quad \sigma_\lambda^- = \epsilon_{\lambda ji} \sigma_{ij}, \quad \lambda, i, j = 1, 2, 3; \quad (10)$$

indexes κ, μ, ν run over $s, \dot{s}$; $\sigma_{\mu\nu} = \frac{i}{2} [\gamma_\mu, \gamma_\nu]$, γ_μ are the Dirac matrices [6]. Hereinafter, we will omit the sign $\pm$. Since the skew-symmetric tensor $\epsilon_{\lambda jk}$ appears in the equation for σ_λ , the generalized Pauli matrixes σ_λ are pseudo-vectors. Therefore, right- and left-helical particles differ from one another.

It follows from the expansion (9) of the wave function $|\psi\rangle$ in a series that the projection P defined by the expression

$$P|\psi\rangle = \frac{1}{2} (|x_\lambda^R\rangle \langle x_\lambda^R| + |x_\lambda^L\rangle \langle x_\lambda^L|) \psi, \quad (11)$$

can be represented in the form

$$P|\psi\rangle = \frac{1}{2} (|x_\lambda^R\rangle \langle x_\lambda^R| + |x_\lambda^L\rangle \langle x_\lambda^L|) \psi = \frac{1}{2} \left\{ \sum_\mu |\xi_\mu\rangle \langle \xi_\mu | \psi \rangle + \sum_\lambda |\xi_\kappa\rangle (\sigma_\lambda)_{\kappa\nu} \langle \xi_\nu | \psi \rangle \right\}. \quad (12)$$

Let us prove that the projector (12) selects states with a given orientation. The right side of the expression (12) in the representation $\{|\xi_i\rangle\}_i$ is written as

$$P\langle \xi_\lambda | \psi \rangle = \frac{1}{2} \left\{ 1 + \sum_\lambda \langle \xi_\lambda | \xi_\kappa \rangle (\sigma_\lambda)_{\kappa\nu} \right\} \langle \xi_\nu | \psi \rangle \quad (13)$$

We can introduce a four-vector $s^\lambda = \langle \xi_\lambda | \xi_\kappa \rangle$ describing a spin of the system. A convolution of the vector s^λ with matrixes σ_λ is assumed on the right side in Eq. (13). It follows from here, that $P = P(s)$ is determined in s -representation by the expression

$$P(s) = \frac{1}{2} (1 + s^\mu \gamma_\mu) \quad s^\mu = \{0, s^\lambda\}. \quad (14)$$

The formula (14) is a known expression for the projection operator that selects bispinors with a given orientation in the rest frame [6]. Hence, it is possible to present the projector P in the form (12). The equality (12) in a matrix form (6) for the representation $\Phi = X = X^R = X^{L\dagger}$ has the following form

$$P|\psi\rangle = \frac{1}{2} (|x_\lambda^R\rangle\langle x_\lambda^R| + |x_\lambda^L\rangle\langle x_\lambda^L|) |\psi\rangle = \frac{1}{2} \left\{ \sum_\mu |\xi_\mu\rangle\langle\xi_\mu| \psi + \sum_\lambda |\xi_\kappa\rangle\langle(\sigma_\lambda)_{\kappa\nu}| \xi_\nu \psi \right\} \quad (15)$$

After left multiplication on $\langle x_\lambda^R|$ the requirement of orthonormal basis $\langle x_{\lambda_i}^R | x_{\lambda_j}^R \rangle = \delta_{ij}$ allows us to rewrite the expression (15) in the form

$$\langle x_\lambda^R | x_\lambda^L \rangle \langle x_\lambda^L | \psi \rangle = \langle x_\lambda^R | \xi_\kappa \rangle (\sigma_\lambda)_{\kappa\nu} \langle \xi_\nu | x_\lambda^L \rangle \langle x_\lambda^L | \psi \rangle \quad (16)$$

Let us introduce the designations

$$x_\lambda = \langle x_\lambda^R | x_\lambda^L \rangle; \quad \xi_\kappa = \langle x_\lambda^R | \xi_\kappa \rangle. \quad (17)$$

Taking into account the expression (17) after simple transformations the expression (16) is written as

$$x_\lambda = \xi_\kappa (\sigma_\lambda)_{\kappa\nu} \xi_\nu. \quad (18)$$

From the above mentioned it follows that to carry out the projection, coordinates should be changed in the wave equation (8) according to Eq. (18) and the wave function ψ should be chosen so that its projection P reads

$$|\psi\rangle = P|\psi\rangle. \quad (19)$$

Let us choose a polar coordinates system $\{\rho_i, \chi_i\}_{i=1}^2$ which obeys $\rho_i = \sqrt{\xi_i^2 + \xi_{i+1}^2}$, $i = 1, 2$, and χ_i is a polar angle in the plane $\{\rho_i, \chi_i\}$, $i = 1, 2$. Then the condition (19) takes the form

$$\langle \rho_1, \rho_2, \chi_1, \chi_2 | \psi \rangle = \langle \rho_1, \rho_2, \chi_1, \chi_2 | P | \psi \rangle = \langle \rho_1, \rho_2, \chi_2 = f(\chi_1) | \psi \rangle, \quad (20)$$

where $f(\chi_1)$ is a function of the polar angle χ_1 . The above-stated allows us to write the equation (8) in the three-dimensional physical space.

Next, we examine a case of the particle with an integer-valued spin in the rest reference frame.

3 A motion equation for a particle with an integer-valued spin

We know that in the rest frame the spin of a system can be described by a three-vector s^μ as

$$s^\mu = \{0, s_1, s_2, s_3\}. \quad (21)$$

Then, the vector $\vec{\xi}_s$ in the equation (8) is determined on ordinary c -numerical space, i.e. space with commutation relations. Since components of the vector $\vec{\xi}_s$ is ordinary c -number, the commutators in the last summand vanish. Hence, we get

$$\left(-\frac{\partial^2}{\partial \xi_s \partial \xi_s} - Z^2 + \omega^2 (\bar{\xi}_s \xi_s) |\Phi\rangle\langle\Phi| \right) |\psi\rangle = 0. \quad (22)$$

Here $\bar{\xi}_s$ is a transposed spinor conjugated to the spinor ξ_s . The obtained equation (22) is the equation for an isotropic harmonic quantum oscillator with the mass $m = \frac{1}{2}$ in two-dimensional space with complex coordinates $\xi_s \stackrel{def}{=} \langle \Phi = X | \xi_s \rangle \equiv \vec{\xi}_s$, $s = 1, 2$ that are components of bispinor in the representation $\Phi = X$, determined on ordinary c -numerical space.

In the next section we will project a trajectory of a particle on three-dimensional space.

4 Transformation to space-time coordinates

Taking into account the condition (21) we calculate the square of module of the expression (18):

$$\begin{aligned} x^\lambda x_\lambda = |x|^2 &= \left| \begin{pmatrix} \xi_s \\ \xi_{\dot{s}} \end{pmatrix}^\dagger \begin{pmatrix} 0 & \sigma_i \\ -\sigma_i & 0 \end{pmatrix} \begin{pmatrix} \xi_s \\ \xi_{\dot{s}} \end{pmatrix} \right|^2 \\ &= |\xi|^2 \begin{pmatrix} \xi_s \\ \xi_{\dot{s}} \end{pmatrix}^\dagger \begin{pmatrix} 0 & \sigma_i^2 \\ \sigma_i^2 & 0 \end{pmatrix} \begin{pmatrix} \xi_s \\ \xi_{\dot{s}} \end{pmatrix} = |\xi|^4. \end{aligned} \quad (23)$$

Here $|\xi|^2 = \sum_{\nu=1}^4 \xi_\nu^2$, the Dirac's representation of matrices γ^μ , $\mu = 0, 1, 2, 3$ was chosen; and it was taken into account that $\sigma_i^2 = 1$, where σ_i are the Pauli matrices. It follows from the obtained expression (23) that

$$\xi_1^2 + \xi_2^2 + \xi_3^2 + \xi_4^2 = r \quad (24)$$

where $r = |x|$. According to the expression (24) the following replacement of variables $\xi_\kappa \rightarrow x_\lambda$ in the motion equation (22) may be performed:

$$\xi_1^2 + \xi_2^2 = \frac{r}{2} + x_3 = u; \quad \xi_3^2 + \xi_4^2 = \frac{r}{2} - x_3 = v \quad (25)$$

As seen from Eq. (25) it is necessary to pass to the polar coordinates $(\rho_i, \chi_i)_{i=1}^2$

$$\rho_i = \sqrt{\xi_i^2 + \xi_{i+1}^2}, \quad \chi_i = \arctan\left(\frac{\xi_{i+1}}{\xi_i}\right), \quad i = 1, 2. \quad (26)$$

In these coordinates the square of an arc element determined by

$$ds_1^2 = d\rho_1^2 + \rho_1^2 d\chi_1^2 = \frac{1}{u} du^2 + u d\chi_1^2; \quad (27)$$

$$ds_2^2 = d\rho_2^2 + \rho_2^2 d\chi_2^2 = \frac{1}{v} dv^2 + v d\chi_2^2 \quad (28)$$

where variables u, v are defined by the expression (25). Evidently, that in the coordinates $\xi_s, \xi_{\dot{s}}$ the equation (22) is solvable by a separation of variables. Therefore, Laplacian can be written in the form $\Delta = \Delta_s + \Delta_{\dot{s}} = \sum_{i=1}^2 \Delta_i$. To get the motion equation in the polar coordinates ρ_i, χ_i , $i = 1, 2$ we use a theorem according to which the Laplacian Δ_i , $i = 1, 2$ in curvilinear orthogonal coordinates q_{1i}, q_{2i} has the form

$$\Delta_i = \frac{1}{h_{1i}h_{2i}} \left\{ \frac{\partial}{\partial q_{1i}} \left(\frac{h_{2i}}{h_{1i}} \frac{\partial}{\partial q_{1i}} \right) + \frac{\partial}{\partial q_{2i}} \left(\frac{h_{1i}}{h_{2i}} \frac{\partial}{\partial q_{2i}} \right) \right\}, \quad (29)$$

where h_{1i} , h_{2i} are the Lamé coefficients in the expression for the square of the differential of arc-length

$$ds_i^2 = h_{1i}^2 dq_{1i}^2 + h_{2i}^2 dq_{2i}^2. \quad (30)$$

By applying this theorem to our case, after the simple transformations we gain the Laplacian in coordinates (u, v, χ_1, χ_2) :

$$\Delta_{u,v,\chi_1,\chi_2} = \left\{ \frac{\partial}{\partial u} \left(u \frac{\partial}{\partial u} \right) + \frac{\partial}{\partial v} \left(v \frac{\partial}{\partial v} \right) + \frac{1}{u} \frac{\partial^2}{\partial \chi_1^2} + \frac{1}{v} \frac{\partial^2}{\partial \chi_2^2} \right\}. \quad (31)$$

Since the degenerate discrete representations of the group $SO(4,2)$ coincide with the projectors defined by subalgebra $\mathfrak{su}(1,1)$ [4], one has a relation $m_1 = m_2 = m$ in the expression for a wave function $\psi = U(u)e^{im_1\chi_1}V(v)e^{im_2\chi_2}$. Therefore the wave function is represented as

$$\psi = U(u)V(v)e^{im\chi}. \quad (32)$$

Here

$$\chi = \chi_2 + \chi_1. \quad (33)$$

Hence, as a manifold $\{u, v, \chi, \eta\} \in \mathbb{R}^4$, $\eta = \chi_2 - \chi_1$, on which motion of the particle is examined, one can choose such a submanifolds being a section

$$\eta = 2\pi a n, \quad n = 1, 2, \dots \quad (34)$$

where a is a constant.

Let us introduce cylindrical coordinates in three-dimensional physical space z, ρ, ϕ :

$$z = x_3 = \frac{u-v}{2}, \quad \rho = \sqrt{uv} \quad \phi = \frac{\chi}{2}. \quad (35)$$

The polar angle ϕ varies within the limits from 0 to 2π , because the angle χ varies from 0 to 4π . Eqs. (34), (35) allows to make the following replacement

$$\frac{\partial}{\partial \chi_2} \left(\frac{\partial}{\partial \chi_1} \right) \Rightarrow \frac{1}{2} \frac{\partial}{\partial \phi}. \quad (36)$$

Substitution of Eqs. (31, 36) into Eq. (22) gives

$$-\left\{\frac{\partial}{\partial u}\left(u\frac{\partial\psi}{\partial u}\right)+\frac{\partial}{\partial v}\left(v\frac{\partial\psi}{\partial v}\right)+\frac{1}{4}\left(\frac{1}{u}+\frac{1}{v}\right)\frac{\partial^2\psi}{\partial\phi^2}\right\}-(Z^2-\omega^2(u+v))\psi=0. \quad (37)$$

Dividing the equation (37) on $u+v$ and multiplying on 4, one gets

$$-\frac{4}{u+v}\left\{\frac{\partial}{\partial u}\left(u\frac{\partial\psi}{\partial u}\right)+\frac{\partial}{\partial v}\left(v\frac{\partial\psi}{\partial v}\right)+\frac{u+v}{4uv}\frac{\partial^2\psi}{\partial\phi^2}\right\}-4\left(\frac{Z^2}{u+v}-\omega^2\right)\psi=0. \quad (38)$$

Denote

$$\xi^2 = \xi_1^2 + \xi_2^2; \quad \eta^2 = \xi_3^2 + \xi_4^2, \quad (39)$$

Because

$$dz^2 + d\rho^2 = (\xi^2 + \eta^2)(d\xi^2 + d\eta^2) = \frac{1}{4}(u+v)\left(\frac{1}{u}du^2 + \frac{1}{v}dv^2\right), \quad (40)$$

the square of arc element

$$ds^2 = dz^2 + d\rho^2 + \rho^2 d\phi^2 \quad (41)$$

in cylindrical 3d-frame is equal to

$$ds^2 = \frac{u+v}{4u}du^2 + \frac{u+v}{4v}dv^2 + uv d\phi^2 \quad (42)$$

in (u,v)-coordinates. Thus, the Laplace operator in 3d-space has the form

$$\Delta_{q_1, q_2, q_3} = \frac{1}{h_1 h_2 h_3} \left\{ \frac{\partial}{\partial q_1} \left(\frac{h_2 h_3}{h_1} \frac{\partial}{\partial q_1} \right) + \frac{\partial}{\partial q_2} \left(\frac{h_3 h_1}{h_2} \frac{\partial}{\partial q_2} \right) + \frac{\partial}{\partial q_3} \left(\frac{h_1 h_2}{h_3} \frac{\partial}{\partial q_3} \right) \right\}, \quad (43)$$

where coefficients h_1, h_2, h_3 determine the square of the arc element as

$$ds^2 = h_1^2 dq_1^2 + h_2^2 dq_2^2 + h_3^2 dq_3^2. \quad (44)$$

From the expression for the Laplace operator

$$\Delta_{u, v, \phi} = \frac{4}{u+v} \left\{ \frac{\partial}{\partial u} \left(u \frac{\partial}{\partial u} \right) + \frac{\partial}{\partial v} \left(v \frac{\partial}{\partial v} \right) + \frac{u+v}{4uv} \frac{\partial^2}{\partial\phi^2} \right\} \quad (45)$$

appearing in Eq. (38) it follows that

$$h_1 = \frac{1}{2}\sqrt{\frac{u+v}{u}}, \quad h_2 = \frac{1}{2}\sqrt{\frac{u+v}{v}}, \quad h_3 = \sqrt{uv}. \quad (46)$$

Comparing expressions (46) and (42) it may be concluded that the Laplacian (45) was written in parabolic 3d-frame u, v :

$$u = R + z, \quad v = R - z, \quad (47)$$

in which the square of the arc element has the form (42). Comparing the expression (25) with the expression (47) one obtains that

$$R = \frac{r}{2}, \quad (48)$$

5 Derivation of wave equations for charged vector-boson on non-compact group

According to expressions (41, 43, 44, 46, 48) it follows that the motion equation (38) in the spherical frame can be rewritten as

$$-\Delta\psi - \frac{2Z^2}{R}\psi = -4\omega^2\psi. \quad (49)$$

After elementary substitution $4\omega^2 \Rightarrow E_1^2 - m_1^2$, $Z^2 \Rightarrow \alpha(E_1 + m_1)$ one gets

$$\left[-\nabla^2 - \left(E_1 + \frac{\alpha}{R} \right)^2 + \left(m_1 - \frac{\alpha}{R} \right)^2 \right] \psi = 0, \quad (50)$$

which, in principle is the Klein-Gordon equation, as it must, with a scalar potential $\varphi(R) = -\imath\alpha/R$ and a vector potential $A(R) = -\imath\alpha/R$, where $A(R)$ is a displacement of the R -th component of the gradient $\vec{\nabla}$. This results from the form of the Laplacian in the spherical coordinates and the facts that $\psi \equiv \{\psi_i\}_{i=1}^3 \stackrel{def}{=} \vec{\psi}$ and the spherical symmetry of a problem yields $\vec{A} \cdot \vec{\nabla}_R \psi \equiv \vec{A} \cdot (\vec{\nabla}_R \times \vec{\psi}) = 0$; where $\vec{\nabla}_R$ is a radial component of gradient $\vec{\nabla}$. Therefore, as a result, the equation (50) can be rewritten as

$$\left[\hat{P}^2 - \left(E_1 + \frac{\alpha}{R} \right)^2 + m_1^2 \right] \psi^* = 0, \quad \hat{P} = \tau_3 \left(P_R + \imath \frac{\mathcal{M}}{R} \right), \quad \mathcal{M} = -\frac{\tau_1}{\sin \theta} p_\phi + \tau_2 p_\theta \quad (51)$$

where ψ^* is the wave function of a particle with an integer spin, which is determined through the generalized spherical function with an integer spin and results from the transformation of the original wave function similar to fermions [7]; the wave function ψ^* is an eigenfunction of the operator $\mathcal{M}$: $\mathcal{M}\psi = m_1\psi$; $\psi = \psi^*/R$; matrices τ_i , $i = 1, 2, 3$ are generators of the rotation group $\text{SO}(3)$ in a matrix form, which correspond to the generalized Pauli matrices for the spherically symmetric Dirac equation; $\hat{P}$ is the relativistic momentum operator obtained by adding components of the vector-potential $\vec{A} = \{A(R), 0, 0\}$ to the ordinary momentum operator which is a derivative (tangential vector): $P_R = p_R + A(R)$, $p_R = -\imath\hbar \frac{\partial}{\partial R}$, $p_\theta = -\imath\hbar \frac{\partial}{\partial \theta}$, $p_\phi = -\imath\hbar \frac{\partial}{\partial \phi}$.

Moreover, it can be seen that when taking into account the explicit expression for eigenvalues Z^2 of the Hamiltonian of two-dimensional harmonic oscillator (22) with the mass $m = \frac{1}{2}$ and eigenfrequency $\omega_0 = 2\omega$:

$$Z^2 = 2\omega(n_r + l + 1), \quad \hbar = 1 \quad (52)$$

then Eq. (49) can be rewritten as

$$-\Delta\psi - \frac{2Z^2}{R}\psi = E_n\psi, \quad (53)$$

where

$$E_n = -\frac{Z^4}{(n_r + l + 1)^2}, \quad m = \frac{1}{2}, \quad \hbar = 1, \quad (54)$$

n_r, l are quantum numbers of the harmonic oscillator with the Laplacian (29). Let us assume that $2Z^2 \Rightarrow e^2$. If e is supposed to be the charge of an electron then the equation (53) is nothing but the equation on eigenfunctions with eigenvalues E_n of a hydrogen-like atom.

6 Asymptotic solution

Apparently, one can write for the relativistic boson (50) that $(E_1^2 - m_1^2)n^2 = \alpha^2(E_1 + m_1)^2$ and $\frac{Z^2}{n^2} = \frac{E_1 - m_1}{\alpha}$. Therefore, the relativistic boson has the following spectrum $E_1 = m_1 \frac{(1 + \frac{\alpha^2}{n^{*2}})}{(1 - \frac{\alpha^2}{n^{*2}})}$ with a mass spectrum $m_1 = \frac{\tilde{Z}^2}{2} \left(1 - \frac{\alpha^2}{n^2} \right)$,

where $n = n_r + l + 1$, $n^* = n_r^* + l^* + 1$, $\tilde{Z}^2 = \frac{Z^2}{\alpha}$. Thus, Eq. (50) describes a compound physical system. Let us choose $\alpha = \imath\gamma$, $\gamma \in \mathbb{R}$, that yields the Coulomb potential. Then, the equation (50) allows relativistic corrections to the spectrum of the hydrogen-like atom in the non-relativistic limit $n(n^*) \rightarrow \infty$ to be found. With the above mentioned consideration, the spectrum in this case has the form

$$E_1 \approx \frac{\tilde{Z}^2}{2} \left(1 - \frac{\gamma^2}{n^{*2}} \right) \left(1 - \frac{\gamma^4}{4n^4} \right) \quad (55)$$

if the spectrum is assumed to consist of two spectral series: odd one $\{n^*\}$ and even one $\{n\} \rightarrow \{2n\}$ and the series are close: $\gamma^2/n^{*2} - \gamma^2/4n^2 = \bar{\epsilon}$, $\bar{\epsilon} \ll 1$. Since the series are close,

$$n^{*2} \approx n^2 + \epsilon, \quad |\epsilon|/n^2 \ll 1. \quad (56)$$

Let us choose ϵ as

$$\epsilon = 2(n - |k|)(\sqrt{k^2 - \gamma^2} - |k|) \approx -(n - |k|)\frac{\gamma^2}{|k|}, \quad (57)$$

where $k = -l, l + 1$.

Introduce designations: $\tilde{Z}^2 \Rightarrow m$, where m is a particle mass. Then, expanding into a series on $\frac{\gamma^2(n-|k|)}{|k|}$, we gain accurate to terms of the order $m\gamma^8$

$$E_1 \approx \frac{m}{2} - \frac{m\gamma^2}{2n^2} - \frac{m\gamma^4}{8n^3} \left(\frac{4}{|k|} - \frac{3}{n} \right) - \frac{m\gamma^6}{8n^4} \left(\frac{3}{n^2} - \frac{8}{n|k|} + \frac{4}{k^2} \right) + O(\gamma^8). \quad (58)$$

From the formula (58) it follows that the first term yields the relativistic rest energy of the compound system considered with a reduced mass $m/2$. The second term yields the Rydberg formula. The third term is a relativistic correction which describes the fine structure of the hydrogen-like atom lines with the fine structure constant $|\gamma|$ in the system of units: $e = 1$, $\hbar = 1$, $c = 1$. The fourth term is a relativistic correction which yields the computational method used.

7 Symmetry of wave equations for a particle with integer-valued spin

Now, let us find symmetry groups of the wave equations (50) and (53). With this goal one defines on bispinor $(\xi_s, \xi_{\dot{s}})^T$ a creation operator $\hat{a}^+$ and an annihilation operator $\hat{a}$ as

$$\hat{a} = \begin{pmatrix} a_s \\ a_{\dot{s}} \end{pmatrix} = \begin{pmatrix} \sqrt{\omega} \left(\xi_s + \frac{1}{\omega} \frac{\partial}{\partial \xi_s} \right) \\ \sqrt{\omega} \left(\xi_{\dot{s}} + \frac{1}{\omega} \frac{\partial}{\partial \xi_s} \right) \end{pmatrix}, \quad (59)$$

$$\hat{a}^+ = \begin{pmatrix} a_s^+ \\ a_{\dot{s}}^+ \end{pmatrix}^T = \begin{pmatrix} \sqrt{\omega} \left(\xi_{\dot{s}} - \frac{1}{\omega} \frac{\partial}{\partial \xi_s} \right) \\ \sqrt{\omega} \left(\xi_s - \frac{1}{\omega} \frac{\partial}{\partial \xi_s} \right) \end{pmatrix}^T \quad (60)$$

where the sign T denotes the transposition operator. Then, we find accurate up to a sign operators commuting with the hamiltonian H from Eq. (22):

$$\begin{aligned} M_\lambda &= (\sigma_\lambda)_{\dot{s}t} a_t a_{\dot{s}}, & M_\lambda^+ &= (\sigma_\lambda)_{\dot{s}t} a_s^+ a_t^+, \\ N_\lambda^a &= (\sigma_\lambda)_{st} a_s^+ a_t, & N_\lambda^b &= (\sigma_\lambda)_{\dot{s}t} a_t^+ a_{\dot{s}}, \\ P &= a_s a_{\dot{s}}, & P^+ &= a_s^+ a_{\dot{s}}^+, \\ H &= 2 + a_s^+ a_s + a_{\dot{s}}^+ a_{\dot{s}}. \end{aligned} \quad (61)$$

Using commutation relations for the annihilation (59) and the creation (60) operators it can be shown that fifteen operators (61) will constitute a basis set of generators for the algebra of a symmetry group similar to the Lie algebra of the group $\text{SO}(4,2)$ or $\text{SO}(6)$. The Lie group $\text{SO}(4,2)$ is non-compact while the Lie group $\text{SO}(6)$ is the compact group. According to "no-go" -theorem [6], [9] taking place for boson degrees of freedom, there are no finite-dimensional unitary representation of the noncompact Lie group. This suggests that any nontrivial union of the Poincaré and an interior symmetry group yields an S matrix which is equal to 1. The equation (50) or (51) describes a particle which wave function is transformed on the infinite-dimensional representation of the noncompact Lie group. From here and from the "no-go" theorem it follows that the obtained relativistic equation (51) has the symmetry of $\text{SO}(4,2)$ and is reasonable for the description in an arbitrary reference frame.

8 Conclusions

The equation (53) is the Schrödinger equation for the hydrogen-like atom. The equation (53) also should describe locally a physical system having a symmetry similar to the noncompact symmetry group $\text{SO}(4,2)$. Fock [8] was first to indicate invariance of the Schrödinger equation for the hydrogen-like atom with respect to four-dimensional rotations. But the equation (53) does not contradict to "no-go" -theorem only in the case if it describes a particle, which wave function is transformed according to representation of a compact group similar to the group $\text{SO}(4,2)$. The group $\text{SO}(4,2)$ is similar to the group $\text{SO}(6)$. The generators of these groups differ by signs of metric multipliers [10]. Hence we conclude, that the equation (53) describes a particle, which wave function is transformed on representation of the compact group $\text{SO}(6)$. This description is acceptable to the rest reference frame when it is possible to neglect the Lorentz boost transformations of the group $\text{SO}(4,2)$.

Thus, within the framework of the projection operator method two wave equations for a vector charged boson are obtained, one of which is precise (equation (50)), and the second equation (the equation (53)) describes a

motion of a vector charged boson approximately. The hydrogen-like atom is the compound quantum system with an integer spin and in the stationary case that does not take into account the Lorentz boost transformations, the replacement of a dynamic symmetry group $SO(4,2)$ on $SO(6)$ yields the non-relativistic Schrödinger equation for the hydrogen-like atom. On this basis, we get the conclusion that the obtained relativistic corrections (58) describe the fine structure of the hydrogen-like atom lines with precision to about γ^8 . In general, infinite-component wave equations have also been examined by Nambu [11] to describe the hydrogen-like atom.

References

- [1] I.P. Volobiuev, Yu.A. Kubishin. *Differential geometry and algebras Lie and their applications in field theory*. Editorial URSS, Moscow, 1998 (in russian).
- [2] L.I. Gurskii, L.I. Komarov, A.M. Solodukhin. Group of Symmetry of the Periodic System of Chemical Elements. *Inter.J.of Quantum Chemistry*, v.72, 499-508 (1999)
- [3] I.D. Feranchuk, L.I. Gurskii, L.I. Komarov, O.M. Lugovskaya, F. Burgäzy and A.Ulyanenko. A new method for calculation of crystal susceptibilities for X-ray diffraction at arbitrary wavelength. *Acta Crystallographica, Section A*, v. 58, 370-384 (2002).
- [4] Barut A.O., Rączka R. *Theory of group representations and applications*. Vol. 1,2. Warszawa. 1977.
- [5] V.Guillemin, S.Sternberg. *Geometric asymptotics*. AMS, Providence, Rhode Island, 1977.
- [6] Michio Kaku. *Quantum field theory*. Oxford. 1993.
- [7] V.A. Fock. *Foundations of quantum mechanics*. Moscow.: "Nauka". 1976 (in Russian).
- [8] V.A. Fock. To the theory of hydrogen atom. *Z.Phys.* v.98, 145 (1935)
- [9] R. Jost. *Helv.Phys.Acta.* **39**, 369 (1966)
- [10] E.B. Aronson, I.A. Malkin, V.I. Manko. Dynamical symmetries in quantum theory. In: *Elementar. partic. and atom kernels.* **5**, 122 (1974) (in Russian).
- [11] Nambu Y. An infinite-component wave equation with hydrogen-like spectrum of mass. *Phys. Rev.* **160**, 1171 (1961)